

AD-A075 919

AEROSPACE CORP EL SEGUNDO CA SPACE SCIENCES LAB

F/G 4/1

A SIMPLE MODEL OF THE MAGNETOSPHERE.(U)

SEP 79 J M LUHMANN , L M FRIESEN

F04701-78-C-0079

UNCLASSIFIED

TR-0079(4960-05)-6

SAMSO-TR-79-95

NL

| OF |
ADA
075919



END
DATE
FILMED
11-79
DDC

LEVEL

(12)
H SE

AD A 075919

A Simple Model of the Magnetosphere

J. G. LUHMANN AND L. M. FRIESEN
Space Sciences Laboratory
Laboratory Operations
The Aerospace Corporation
El Segundo, Calif. 90245

12 September 1979

Interim Report

APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED

DDC
RECEIVED
NOV 1 1979
A

DDC FILE COPY


Prepared for
SPACE AND MISSILE SYSTEMS ORGANIZATION
AIR FORCE SYSTEMS COMMAND
Los Angeles Air Force Station
P.O. Box 92960, Worldway Postal Center
Los Angeles, Calif. 90009

79 11 01 008

This interim report was submitted by The Aerospace Corporation, El Segundo, CA 90245, under Contract No. F04701-78-C-0079 with the Space and Missile Systems Organization, Contracts Management Office, P.O. Box 92960, Worldway Postal Center, Los Angeles, CA 90009. It was reviewed and approved for The Aerospace Corporation by G. A. Paulikas, Director, Space Sciences Laboratory. Gerhard E. Aichinger was the project officer for Mission-Oriented Investigation and Experimentation (MOIE) Programs.

This report has been reviewed by the Information Office (OI) and is releasable to the National Technical Information Service (NTIS). At NTIS, it will be available to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication. Publication of this report does not constitute Air Force approval of the report's findings or conclusions. It is published only for the exchange and stimulation of ideas.


Gerhard E. Aichinger
Project Officer

FOR THE COMMANDER


Frank J. Bane, Chief
Contracts Management Office

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

19 REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
18 SAMS0-TR-79-95		9
4. TITLE (and Subtitle)	5. TYPE OF REPORT & PERIOD COVERED	
6 A SIMPLE MODEL OF THE MAGNETOSPHERE	Interim rept.	
7. AUTHOR(s)	8. PERFORMING ORG. REPORT NUMBER	9. CONTRACT OR GRANT NUMBER(s)
10 Janet M. /Luhmann and Lynn M. /Friesen	14 TR-0079(4960-05)-6	15 F04701-78-C-0079
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
The Aerospace Corporation El Segundo, California 90245		12 24
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE
Space and Missile Systems Organization Los Angeles Air Force Station Los Angeles, California 90009		11 12 September 1979
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES
		21
		15. SECURITY CLASS. (of this report)
		Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)		
Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
Magnetosphere model		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
<p>A phenomenological magnetic field model for the earth's magnetosphere is constructed from a dipole field and a uniform field directed sunward in the northern hemisphere and antisunward in the southern hemisphere. The properties of this simple model are compared with those of several other quantitative models. The present model is found to be more suitable for calculations than some other simple models in cases where the distant ($>13 R_e$) magnetotail configuration is important. Moreover, this model is easily adaptable to changes in the field geometry and to the description of magnetotail asymmetries.</p>		

DD FORM 1473
(FACSIMILE)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

407 512

JOB

PREFACE

The authors would like to thank Michael Schulz for many helpful comments relating to the presentation of this model, especially with respect to the method of smoothing the equatorial discontinuity. The assistance of Henry H. Hilton in the programming effort involved in the field line tracing is also gratefully acknowledged.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or special
A	

CONTENTS

PREFACE	1
INTRODUCTION	5
THE BASIC MODEL	5
PROPERTIES OF THE BASIC MODEL	7
TILTED DIPOLE	14
ADDITION OF ROTATION	16
CONCLUDING REMARKS	19
REFERENCES	21

FIGURES

1.	The basic magnetospheric model described by eq. (1)	8
2.	Field lines described by eq. (1) in the noon-midnight plane, drawn for colatitudes within 50° of the poles at intervals of 5°	9
3.	Comparison of field strengths along field lines originating at various latitudes in the present model (eq. (1)) and the 13-term Mead model	10
4.	Comparison of field lines originating at various magnetic latitudes Λ in the Olson and Pfitzer (1974) (OP 74), Mead and Fairfield (1975) (MF 75) and present magnetospheric field models in the noon-midnight plane	12
5.	Selected field lines in the noon-midnight plane of the two term (2T) and six term (6T) field models (Williams and Mead, 1965) which are used in the calculation of cosmic ray trajectories in the magnetosphere	13
6.	The magnetosphere model described by eq. (5)	15
7.	The tilted magnetosphere model described by eq. (6) with $\alpha = 30^\circ$ as it appears for the same field lines as shown in Fig. 1	17
8.	Illustration of the effect of adding the "rotation" term $-K/r \sin \theta$ to \vec{B}	18

INTRODUCTION

Models of the magnetic field within the magnetosphere have been developed for the purpose of numerical analyses such as particle trajectory tracing and the ordering of satellite data. Roederer (1969) reviewed some of the early models. Within the last few years, Olson and Pfitzer (1974), Choe and Beard (1974) and Mead and Fairfield (1975) have developed sophisticated multiple-term models which include magnetospheric currents and the tilt of the earth's dipole axis with respect to the incident solar wind direction. For many purposes, however, simple models are desirable because of their mathematical tractability. Some simplified models have been formulated from truncated versions of the series expansions describing the sophisticated models (e.g. Williams and Mead, 1965, Olson and Pfitzer, 1974).

In this report an alternative simple model is described. The basic model is constructed from a dipole field and a uniform field directed sunward in the northern hemisphere and antisunward in the southern hemisphere. Some properties of the model are discussed and compared with several other quantitative models. The incorporation of tilt or rotation-like features by the addition of terms is demonstrated to illustrate the flexibility of the model.

THE BASIC MODEL

The magnetic field of the present untilted magnetospheric model is described by the equation

$$\bar{B} = \mu \bar{\nabla} (\cos \theta / r^2) + B_T \sin \theta \cos \phi \hat{x} \quad (1)$$

$$B_T > 0 \quad 0 < \theta < \pi/2$$

$$B_T < 0 \quad \pi/2 < \theta < \pi$$

in which θ is colatitude, ϕ is magnetic longitude measured from midnight, r is the geocentric distance and \hat{x} is the unit vector directed perpendicular to the dipole axis. The first term in (1) describes the dipole field, and the second term describes a current-sheet field which points in opposite directions in the northern and southern hemispheres. The current sheet extends throughout the magnetosphere, effectively replacing the magnetopause and ring currents in the inner magnetosphere. Two constants, μ and B_T , describe the dipole field and the current sheet field, respectively. At large distances from the dipole, the field in the magnetotail is approximately equal to B_T .

The components of the magnetic field given by eq. (1) are:

$$B_r = -2\mu \cos \theta / r^3 + B_T \sin \theta \cos \phi \quad (2)$$

$$B_\theta = -\mu \sin \theta / r^3 + B_T \cos \theta \cos \phi$$

$$B_\phi = -B_T \sin \phi$$

where B_T changes sign at the equatorial plane as specified above.

PROPERTIES OF THE BASIC MODEL

The divergence of the field given by eq. (1) vanishes, as it must. It was found that the value $B_T = .00015$ produces a magnetosphere which resembles other quantitative models when the dipole field strength $\mu = .31$ is invoked and when r is measured in earth radii. Several projected views of the 3-dimensional model obtained with these parameters are shown in Fig. 1.

Figure 2 shows the field lines in the noon-midnight plane only. The dimpled magnetopause of this model is located at $r \approx 10$. The colatitude θ_c and radial distance r_c of the noon cusp, where the field lines separate to form the tail, can be found by setting \bar{B} (eq. (2)) equal to zero. The cusp occurs at the constant colatitude

$$\theta_c = \tan^{-1} \sqrt{2} = 54.7^\circ \quad (3)$$

For the parameters $\mu = .31$ and $B_T = .00015$ selected above, the $\bar{B} = 0$ point is located at the radial distance

$$r_c = [\mu \tan \theta_c / B_T]^{1/3} = 14.3 \quad (4)$$

The magnitude of \bar{B} along field lines originating at various dipole latitudes is shown in Fig. 3 together with similar data given by Mead (1964) for his 13-term model. Although the minimum \bar{B} points in the present model lie equatorward of the minima in the Mead



Fig. 1. The basic magnetospheric model described by eq. (1). Projected views from above the equatorial plane (top), in the noon-midnight plane (center), and from the sun (bottom) are shown. Field lines are drawn for colatitudes within 50° of the poles at intervals of 5° and for longitude intervals of 18° .




Fig. 2. Field lines described by eq. (1) in the noon-midnight plane, drawn for colatitudes within 50° of the poles at intervals of 5°




Fig. 3. Comparison of field strengths along field lines originating at various latitudes in the present model (eq. (1)) and the 13-term Mead model

model, the behavior of \bar{B} along a field line is similar. A qualitative comparison with several highly sophisticated models based on spacecraft magnetometer data is illustrated in Fig. 4, which shows selected field lines in the noon-midnight plane. Aside from the dimple in the field lines near the dayside magnetopause, the present simple model appears to agree with the sophisticated models about as well as the sophisticated models agree with each other (e. g. see Walker, 1976). However, the diurnal variation at constant r that occurs in these other models and in the observed field does not occur in the model described by eq. (1).

Selected field lines of simplified models that are employed in particle trajectory calculations for cosmic ray access studies (Gall and Orozco, 1974) are illustrated in Fig. 5. One obvious difference between these models and the present model is the configuration of the magnetotail. In the present model the magnetotail is composed of quasi-parallel field lines that gradually converge toward the equatorial plane where they reverse direction. Many of the other simple models are actually intended for use only at $r \lesssim 13$ earth radii because they have tail field lines that diverge or show other unrealistic behavior (e. g. see Olson and Pfitzer, 1974). Since the present model more accurately represents the observed magnetotail (e. g. see Willis and Pratt 1972), it is superior for simulating effects related to the tail configuration.




Fig. 4. Comparison of field lines originating at various magnetic latitudes Λ in the Olson and Pfitzer (1974) (OP 74), Mead and Fairfield (1975) (MF 75), and present magnetospheric field models in the noon-midnight plane




Fig. 5. Selected field lines in the noon-midnight plane of the two term (2T) and six term (6T) field models (Williams and Mead, 1965) which are used in the calculation of cosmic ray trajectories in the magnetosphere

A primary shortcoming of the present model is the discontinuity of \bar{B} in the equatorial plane. However, this discontinuity is easily smoothed in numerical analyses by introducing a factor $\tanh (r \cos \theta / \delta)$, where δ is a constant, in the tail field term:

$$\bar{B} = \mu \bar{\nabla} (\cos \theta / r^2) + B_T \sin \theta \cos \phi \tanh (r \cos \theta / \delta) \hat{x} \quad (5)$$

(M. Schulz, personal communication). The effect of this modification, as shown by Fig. 6, is essentially that of the addition of a current sheet of "thickness" δ in the equatorial plane. The value of δ determines the colatitude of the last "closed" field lines and the shape of the magnetopause near the equator.

TILTED DIPOLE

The dipole in the present model can be tilted with respect to the tail by rotating the coordinate system for the current-sheet field:

$$B_r = -2\mu \cos \theta / r^3 + B_T (\sin \theta \cos \phi \cos \alpha - \cos \theta \sin \alpha)$$

$$B_\theta = -\mu \sin \theta / r^3 + B_T (\cos \theta \cos \phi \cos \alpha + \sin \theta \sin \alpha)$$

$$B_\phi = -B_T \sin \phi \cos \alpha$$

(6)

$$B_T > 0 \quad \cos \theta > -\sin \theta \cos \phi \tan \alpha$$

$$B_T < 0 \quad \cos \theta < -\sin \theta \cos \phi \tan \alpha$$




Fig. 6. The magnetosphere model described by eq. (5). Field lines in the noon-midnight plane are shown for two values of δ

where α is the angle between the dipole axis and the tail field direction \hat{x} . Selected field lines for the tilted model with $\alpha = 30^\circ$ are shown in Fig. 7. As in other models with a tilted dipole (e.g., Mead and Fairfield, 1975), some field lines appear to penetrate the magnetopause. These lines could be eliminated if the appropriate magnetopause currents were included.

ADDITION OF ROTATION

A rotation-like effect can be incorporated in the basic model (2) by adding a term to B_ϕ which describes a cylindrical vortex oriented along the dipole axis:

$$B_\phi = -B_T \sin \phi - K/r \sin \theta \quad (7)$$

Here K characterizes the vortex strength, or rotational velocity. Although this is not a particularly realistic representation of the consequences of the rotation of the magnetosphere, it is qualitatively consistent with the behavior that may be found at latitudes above the region of corotation.

Figure 8 illustrates that the effect of the above addition to B_ϕ is an asymmetric distortion of the field lines and the magnetopause. In fact, a local-time asymmetry in the magnetopause shape, similar to the distortion produced by the rotation term in eq. (8), has been observed (Fairfield and Mead, 1975, McDiarmid, et. al., 1976).





Fig. 7. The tilted magnetosphere model described by eq. (6) with $\alpha \approx 30^\circ$ as it appears for the same field lines as shown in Fig. 1



$K = .06$



$K = .30$

Fig. 8. Illustration of the effect of adding the "rotation" term $-K/r \sin \theta$ to \vec{B} . Equatorial plane projections of the field lines used in Fig. 1 are shown for two values of K

CONCLUDING REMARKS

It is noteworthy in the present context that Mead and Fairfield (1975), in their report describing a 17-term model obtained by fitting spacecraft magnetometer data, found that the predominant non-dipole term in the series expansion of \bar{B} described a field directed sunward in the northern hemisphere and antisunward in the southern hemisphere.

The advantages of using the simple model that has been described here are obvious. The step-wise calculation of particle trajectories in a model of the magnetospheric magnetic field with a realistic magnetotail can be carried out using minimal computer time. Moreover, the effects on the magnetosphere of the spatial nonuniformity and temporal variability of the solar wind are simply modeled by adjusting B_T . For example, asymmetric distortions can be described by $B_T = B_T(x, y, z)$. The possibilities afforded by the parameterization of B_T , together with the tilt and rotation options, make this an unusually flexible magnetospheric model for both qualitative and quantitative analyses.

REFERENCES

- Choe, J. Y. and D. B. Beard, The compressed geomagnetic field as a function of dipole tilt, *Planet. Space Sci.* 22, 595, 1974.
- Fairfield, D. H. and G. D. Mead, Magnetospheric mapping with a quantitative geomagnetic field model, *J. Geophys. Res.* 80, 535, 1975.
- Gall, R. and A. Orozco, Review on the usage of magnetospheric models, *J. Geophys. Res.* 79, 293, 1974.
- McDiarmid, I. B., E. E. Budzinski and J. R. Burrows, Comparison of the Mead-Fairfield magnetic field model with particle measurements, *J. Geophys. Res.* 81, 3459, 1976.
- Mead, G. D., Deformation of the geomagnetic field by the solar wind, *J. Geophys. Res.* 69, 1181, 1964.
- Mead, G. D. and D. H. Fairfield, A quantitative magnetospheric model derived from spacecraft magnetometer data, *J. Geophys. Res.* 80, 523, 1975.
- Olson, W. P. and K. A. Pfitzer, A quantitative model of the magnetospheric magnetic field, *J. Geophys. Res.* 79, 3739, 1974.
- Roederer, J. G., Quantitative models of the magnetosphere, *Rev. Geophys. Space Phys.* 7, 77, 1969.

Walker, R. J., An evaluation of recent quantitative magnetospheric magnetic field models, *Rev. Geophys. and Space Phys.* 14, 411, 1976.

Williams, D. J. and G. D. Mead, Nightside magnetosphere configuration as obtained from trapped electrons at 1100 kilometers, *J. Geophys. Res.* 70, 3017, 1965.

Willis, D. M. and R. J. Pratt, A quantitative model of the geomagnetic tail, *J. Atmos. Terr. Phys.* 34, 1955, 1972.

LABORATORY OPERATIONS

The Laboratory Operations of The Aerospace Corporation is conducting experimental and theoretical investigations necessary for the evaluation and application of scientific advances to new military concepts and systems. Versatility and flexibility have been developed to a high degree by the laboratory personnel in dealing with the many problems encountered in the nation's rapidly developing space and missile systems. Expertise in the latest scientific developments is vital to the accomplishment of tasks related to these problems. The laboratories that contribute to this research are:

Aerophysics Laboratory: Launch and reentry aerodynamics, heat transfer, reentry physics, chemical kinetics, structural mechanics, flight dynamics, atmospheric pollution, and high-power gas lasers.

Chemistry and Physics Laboratory: Atmospheric reactions and atmospheric optics, chemical reactions in polluted atmospheres, chemical reactions of excited species in rocket plumes, chemical thermodynamics, plasma and laser-induced reactions, laser chemistry, propulsion chemistry, space vacuum and radiation effects on materials, lubrication and surface phenomena, photo-sensitive materials and sensors, high precision laser ranging, and the application of physics and chemistry to problems of law enforcement and biomedicine.

Electronics Research Laboratory: Electromagnetic theory, devices, and propagation phenomena, including plasma electromagnetics; quantum electronics, lasers, and electro-optics; communication sciences, applied electronics, semiconducting, superconducting, and crystal device physics, optical and acoustical imaging; atmospheric pollution; millimeter wave and far-infrared technology.

Materials Sciences Laboratory: Development of new materials; metal matrix composites and new forms of carbon; test and evaluation of graphite and ceramics in reentry; spacecraft materials and electronic components in nuclear weapons environment; application of fracture mechanics to stress corrosion and fatigue-induced fractures in structural metals.

Space Sciences Laboratory: Atmospheric and ionospheric physics, radiation from the atmosphere, density and composition of the atmosphere, aurorae and airglow; magnetospheric physics, cosmic rays, generation and propagation of plasma waves in the magnetosphere; solar physics, studies of solar magnetic fields; space astronomy, x-ray astronomy; the effects of nuclear explosions, magnetic storms, and solar activity on the earth's atmosphere, ionosphere, and magnetosphere; the effects of optical, electromagnetic, and particulate radiations in space on space systems.

THE AEROSPACE CORPORATION
El Segundo, California